ACTH PROTOCOL FOR OMS

Patient Name:	Ht (cm)	S.A Acthar @ 80 IU/C0
Date:	Wt (kg)	Dose = (IU x SA)
Date of Birth:		80

Week	IU per m²	Frequency	Dose=CC
1	75	BID	
2	75	QD	
3	75	QOD	
4	75	QOD	
5	70	QOD	
6	70	QOD	
7	70	QOD	
8	70	QOD	
9	65	QOD	
10	65	QOD	
11	55	QOD	
12	55	QOD	
13	50	QOD	
14	50	QOD	
15	45	QOD	
16	45	QOD	
17	40	QOD	
18	40	QOD	
19	40	QOD	
20	40	QOD	
21	35	QOD	
22	35	QOD	
23	35	QOD	
	35	QOD	
24 25	30	QOD	
26	30	QOD	
27	30	QOD	
28	30	QOD	
29	25	QOD	
30	25 25	QOD	
31	25 25	QOD	
32	25 25	QOD	
33	20	GOD	
34		QOD	
3 4 35	20	QOD	
	20 20	QOD	
<u>36</u> <u>37</u>	20 18	QOD	
38	18	QOD	
39		QOD	
40	18		
41	18 16	QOD	
41	16	QOD	
43		QOD	
44	14	QOD	
45	14 12	QOD	
		QOD	
46	12	QOD	
47	10	QOD	
48	10	QOD	
49	8	QOD	
50	8	QOD	
51	6	QOD	
52	6	QOD	

BID = twice daily QD = once daily QOD = every other day

The National Pediatric Myoclonus Center disclaims all liability in connection with the use of medical information contained in this protocol and website. Important decisions about treatment must be made by individuals with their health care providers.

Michael R. Pranzatelli, MD